

Short Communication

Flattened Nose Revealing Syphilitic Aortitis

Zakia Douhi^{1*}, Imane Couissi¹, Hajar El Bennaye¹, Hanane Baybay¹, Fatima Zahra Mernissi¹

¹Department of Dermatology, University Hospital Hassan II, Morocco

*Correspondence author: Zakia Douhi, Department of Dermatology, University Hospital Hassan II, Fes, Morocco; Email: drzakia1@hotmail.com

Citation: Douhi Z, et al. Flattened Nose Revealing Syphilitic Aortitis. J Dermatol Res. 2023;4(1):1-3.

<https://doi.org/10.46889/JDR.2023.4107>

Received Date: 23-03-2023

Accepted Date: 18-04-2023

Published Date: 24-04-2023



Copyright: © 2023 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license

(<https://creativecommons.org/licenses/by/4.0/>).

Abstract

Flattened nose also called syphilis nose is a deformation of the nose due to the destruction of the cartilage of the nasal septum: the lower segment of the nose collapses, undergoes recoil and retract, becoming embedded in the upper segment. It is characteristic of congenital and tertiary syphilis. We report the case of syphilitic aortitis revealed by the flattened appearance of the nose.

Keywords: Flattened Nose; Tertiary Syphilis; Aortitis

Introduction

Syphilis, before the era of antibiotics, reigned among the leading causes of cardiovascular disease [1]. In its tertiary phase, it can be responsible for several vascular manifestations of varying severity. It occurs in 30% of patients not treated during the primary phase, symptomatic on 10 to 15% of cases [2]. We illustrate through this observation a case of syphilitic aortitis revealed by the flattened appearance of the nose.

Observation

A 45-year-old man, with no particular medical history, was seen in dermatological consultation for necrotic lesion on the lower limb. Clinic examination found necrotic plaques on the left leg and foot with bilateral edema and weak peripheral pulses (Fig. 1). The rest of the examination showed a collapse of the root of the nose (Fig. 2), not noted by the patient and not objectified in his old photos, giving the appearance of a lorgnette nose. Tertiary syphilis was suspected and confirmed by syphilitic serology (VDRL 1/16 TPHA > 2560). The evolution was marked by an extension of the lesions taking all the leg and the left foot, surmounted by place by hemorrhagic

and meliceric crusts. The vascular examination found an abolition of the tibial and femoral pulses. An abdominal CT angiography was performed objectifying a chronic aorto bi-iliac occlusion (Fig. 3), for which he benefited from a trans-femoral amputation and was put on statin, antiplatelet, preventive anticoagulation and penicillin.



Figure 1: Necrotic plaques on the left leg and foot with bilateral edema.



Figure 2: Lorgnette nose.




Figure 3: Angiography aorto bi-iliac occlusion.

Discussion

Syphilis is a sexually transmitted and contagious infectious disease caused by *Treponema pallidum*. We distinguish currently recent syphilis grouping together primary-secondary syphilis and latent syphilis of less than one year for Anglo-Saxons or two years for Americans and late syphilis grouping together latent syphilis over a year or two years and syphilis tertiary [3]. Cutaneous manifestations of tertiary syphilis are diverse, dominated by gums, chronic ulcerations and tubercles. However, the most cited cardiovascular manifestations in the literature are: aortic aneurysm, aortic valvular insufficiency, aortic root dilatation and coronary ostial stenosis. Aortic occlusion has been rarely reported in the literature, but remains very noisy in terms of clinical. Although often regarded as unexpected diagnosis, actually cases of cardiovascular syphilis continue to be reported. Presumably, *Treponema pallidum* invades the aortic wall and the inflammatory response progresses towards obliterative endarteritis and necrosis of the muscular and the elastic fibers in the aortic media. The consequent is the weakness of the aortic wall, which can lead to severe complications [4].

Conclusion

Even in the absence of cardiovascular risk factors or primary infection, cardiovascular syphilis has to stay in mind, specially in young sexually active patients. It should not be forgotten the « great imitator » character of the syphilis.

Conflict of Interest

The authors have no conflict of interest to declare.

References

1. A Mudry. Nose like the foot of a pot. Eur Ann Otorhinolaryngol Head Neck Dis. 2021;138(3):205-7.
2. Jackman Jr JD, Radolf JD. Cardiovascular syphilis. Am J Med. 1989;87:425-33.
3. Saraiva RS, Cesar CA, de Mello MAA, Bras R. Syphilitic aortitis: Diagnosis treatment. Case Report. Cir Cardiovasc. 2010;25:415-8.
4. Drago F, Merlo G, Rebora A, Parodi A. Syphilitic aortitis and its complications in the modern era. G Ital Dermatol Venereol. 2018;153(5):698-706.

Journal of Dermatology Research



Publish your work in this journal

Journal of Dermatology Research is an international, peer-reviewed, open access journal publishing original research, reports, editorials, reviews and commentaries. All aspects of dermatological health maintenance, preventative measures and disease treatment interventions are addressed within the journal. Dermatologists and other researchers are invited to submit their work in the journal. The manuscript submission system is online and journal follows a fair peer-review practices.

Submit your manuscript here: <https://athenaeumpub.com/submit-manuscript/>