

No Impact of Psychosocial Stress on Glioblastoma Development

Durrechou Q¹, Engelhardt J², Jecko V³, Planchon C², Larroquette M¹, Bronnimann C^{1*}

¹Oncology Unit, Hopital Saint André, 1 Rue Jean Burguet 33000 Bordeaux, France

²Neurosurgery B Department, Hôpital Pellegrin, 1 Pace Amélie Raba Léon 33000 Bordeaux, France

³Neurosurgery A Department, Hôpital Pellegrin, 1 Pace Amélie Raba Léon 33000 Bordeaux, France

*Correspondence author: Charlotte Bronnimann, Oncology Unit, Hopital Saint André, 1 Rue Jean Burguet 33000 Bordeaux, France;

Email: charlotte.bronnimann@chu-bordeaux.fr

Abstract

Citation: Durrechou Q, et al. No Impact of Psychosocial Stress on Glioblastoma Development. *J Neuro Onco Res.* 2026;6(1):1-8.

<https://doi.org/10.46889/JNOR.2026.6105>

Received Date: 21-11-2025

Accepted Date: 18-01-2026

Published Date: 25-01-2026



Copyright: © 2026 The Authors. Published by Athenaeum Scientific Publishers.

This is an open access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

License URL:

<https://creativecommons.org/licenses/by/4.0/>

glioblastoma or survival.

Purpose: Psychosocial stress represents an important source of questions in the potential implication of origin of cancer. The aim of the study was to assess stress prevalence prior to diagnosis of wild-type IDH glioblastoma.

Methods: This prospective single-center study enrolled consecutive new cases of wild-type IDH glioblastoma diagnosed between December 2019 and March 2021 at the University Hospital of Bordeaux. A standardized patient self-assessment stress questionnaire explored both the presence of stressor exposure and the intensity of patient stress level prior to the diagnosis. Four groups were included: high stressors/high stress, high stressors/low stress, low stressors/low stress and low stressors/high stress. Patient characteristics were collected. Statistical analysis was based on the Chi-square test and the Kaplan-Meier survival estimator.

Results: Sixty-four patients with a median age of 66 years were included. Glioblastoma involved predominantly the frontal lobe (39%). Thirty-six patients (56.3%) presented a low stressor/low stress profile. Stress corresponded mainly to the death of a loved one or to family health problems. Among working professionals, 20 patients (67.5%) reported low-intensity work stress. A history of depression was found in 30%. Progression-free survival at 6 months was 45.3% and median overall survival was estimated to be 16.5 months. Level and presence of stress did not differ based on location of tumour. No association was found between stress and tumour progression or overall survival.

Conclusion: A majority of patients in this study had low exposure to stressors as well as low stress level. Psychological stress did not seem to favour the emergence of

Keywords: Psychosocial Stress; Glioblastoma; Stressors; Stress Level

Introduction

Stress corresponds to the psychological, physical and emotional reaction that a subject can develop to cope with situations and pressures imposed on them [1]. By activating the adrenergic system and the cortisol pathway, stress becomes responsible for physiological changes (increase in blood pressure, acceleration of the heart rate, disruption of circadian rhythms, etc.) and the hypothesis of an association between stress and increased health risk emerged many years ago [2]. The impact of stress on increased risk of morbidity and mortality is now well established, particularly with cardiovascular diseases. It includes different types of stressful events (bereavement, work, trauma, financial problems, etc) and can be spread over a very variable time scale [3-6]. In oncology, the implication of stress on oncogenesis and tumour progression remains a source of questions. A strong association between stress and cancer has not clearly been established yet [7-9]. In primary central nervous tumours, few risk factors have been identified [10,11]. The potential presence of an identified psychosocial stress prior to diagnosis represents an

important source of routine questions for both patients and caregivers. To our knowledge, no study has yet evaluated the impact of acute or chronic stress on the incidence of brain tumours. The study aimed to assess prospectively the prevalence of exposure to acute and chronic psychosocial stress and the response of patient to the stress in a consecutive cohort of newly diagnosed patients with wild-type IDH glioblastoma at the University Hospital of Bordeaux, France using a standardized self-assessment scale [12]. Secondary objectives were to analyze the potential association between stress and glioblastoma progression and survival.

Methodology

The study included prospectively all adult patients diagnosed for a newly wild-type IDH glioblastoma from December 2019 to March 2021 at the University Hospital of Bordeaux. Exclusion criteria were the presence of a mutant IDH glioblastoma and the presence of advanced cognitive disorders incompatible with filling in the questionnaire. Due to the Covid-19 health crisis during 2020 (confinement, teleworking, telephone announcement) and some cases of intense psychological suffering at the announcement of the diagnosis, some eligible patients could not complete the questionnaire and were therefore not included in the present study. All patients included signed an informed consent form. We collected patient characteristics included clinical, radiological, oncological management and follow-up. Habits reported included excessive alcohol consumption or smoking over 10 pack-years. A history of depression corresponded to an active or past depression for major depression of DSM criteria. Multifocal location corresponded to a presentation with at least 2 consecutive lobes involved with tumoral contiguity. Tumour characteristics included neuropathology according to the 2016 WHO classification of brain tumours and methylation status of the MGMT promoter. Time of first symptoms, time of diagnosis corresponding to surgery and time of acute stress, if present, were reported. At the time of the announcement of the glioblastoma diagnosis, the patient was included and received the validated stress questionnaire from the specialist nurse or the neuro-oncologist separated into two different parts. The first part of the questionnaire entitled "Brief Rating Scale for Stressors" includes 8 questions and the answers to each question are calibrated with a score from 1 to 6 (from "no" to "extremely"). The patient then related the potential history of psychological trauma experienced (any time in their life) as well as their exposure to stressors (health, family, loss, sentimental, financial and professional or other stressors). If recent trauma was identified in the two years prior to diagnosis, the patient was asked to provide written details of this trauma. A global score was calculated and it was then possible to define a stressor intensity level (high level of stressor exposure if score above 18). Question 8 of this part (Do I have a disease? Rate its severity and the discomfort for you) was analyzed independently from the rest of the questionnaire and new intervals were recalculated. The second part of the questionnaire entitled "Brief stress rating scale" included 11 questions about how the stress was expressed through character traits, emotivity, potential psychosomatic manifestations as well as feelings about their health. After adding points obtained for each question, a global score was calculated to define a stress intensity level (high level of stress if score above 30). Patients were separated into 4 different categories according to their level of stressor exposure and stress response (category 1: high stressors /high stress; category 2: high stressors / low stress; category 3: low stressors / low stress; category 4: low stressors / high stress). Patient characteristics as well as Progression Free Survival (PFS) and Overall Survival (OS) were analyzed using a Chi-square test. The Kaplan Meier estimator and the Log Rank test enabled the development and comparison of survival curves. Depending on calculated stressor subgroups, we explored eventual association with tumour location, history of depression, beta-blocker management, Overall Survival (OS) and Progression Free Survival (PFS) with Chi-square test.

Results

65 patients were included in this study with a median age at diagnosis 66 years. One patient was excluded due to IDH mutant status. Patients characteristics are presented in Table 1. Glioblastoma mainly involved the frontal lobe (39%). The methylation status of MGMT promoter was found in 20 cases among the 50 cases assessed. Patients with professional activity at diagnosis represented 30 patients (47%) of the cohort. A history of chronic disease at diagnosis was reported in 46 patients with a majority of high blood pressure in 26 patients, dyslipidemia in 17 patients, diabetes in 11 patients and coronary disease in 10 patients. A history of depression was reported for 19 patients (30%) and 12 patients (19%) were taking psychotropic medication at the time of diagnosis (7 patients with anxiolytic treatment such as benzodiazepines and 5 with antidepressant treatment). 11 patients (17%) of the cohort were taking a beta-blocker treatment for cardiovascular disease at diagnosis. Clinical presentation was mainly a neurological deficit in 33 patients and performance status was less or equal to 1 in 45 patients (70%). Thirty patients (47%) underwent a brain biopsy while 34 patients (53%) had a surgical resection including 9 (26%) with complete resection and 25 (74%) subtotal resection. Four patients with complete resection had a Carmustine wafer implantation during surgery. After a multidisciplinary meeting, a concomitant radio-chemotherapy was the oncological treatment proposed for 57 patients (89%).

Treatment with Temodal alone was decided for 6 patients (9%). One patient underwent palliative care after surgery.

After category recombination, 36 patients (56.3%) presented a "low stressor / low stress" profile (Table 1). According to the first question regarding the experience of psychological trauma, 33 patients (52%) reported low or absence of exposure to psychological trauma (score 1 to 3). The loss of a loved one accounted for 80% of the psychological traumas described by patients. Forty patients (63%) reported low or no acute stress in the 2 years prior to diagnosis and the existence of extreme acute stress was found in only 10 patients (15.6%). Family health problems were the most represented category of stressful events (31%) followed by the loss of a family member (27.5%) and job strain (27.5%). Divorces and other unclassified stressful events each accounted for 7% of stressful events. Among all patients, 37 patients (57.8%) did not mention any job strain and among the 30 working patients, 20 (67.5%) reported low job strain. After analyzing all the results of the first part of the questionnaire, a large majority of patients (67%) had a low or very low level of stressors. Regarding question 8 of the first part of the questionnaire, 39 patients (62.9%) considered the diagnosis of glioblastoma with a significant or extreme severity. Similarly, the second part of the questionnaire showed that 44 patients (69%) had a low or very low level of stress.

For patients on beta-blocker medication, the stress level was low or very low in 73% of cases but the difference was not significant ($p=0,754$). Patients with frontal tumour location had a non-significant increase in stress level (40% vs 21%; $p= 0,121$) and depression rate did not appear to be affected by tumour location ($p = 0,973$). The multifocal nature of the tumour was the only parameter that significantly impacted progression ($p = 0,015$) and survival ($p = 0,002$) negatively. There was no significant difference in stress between right and left hemisphere tumour, nor between dominant and non-dominant hemisphere ($p=0,450$). Furthermore, there appeared to be no significant difference between men and women regarding the level of exposure to stressors (including job strain) ($p = 0,184$) or the level of stress ($p=0,112$). Based on these results, exposure to stressors as well as the intensity of the stress level appear to be relatively low in the glioblastoma population.

Median time between reported stress and diagnosis was 11 months (Fig. 1). Median overall survival was 16,5 months and 6-months progression free survival was 45,3%. No significant association was demonstrated between tumour progression, overall survival and exposure to stressors or stress level. History of depression ($p = 0,755$), use of beta-blockers ($p = 0,683$) and MGMT status ($p = 0,309$) did not appear to significantly impact overall survival.




Figure 1: Time between stress - symptom - glioblastoma diagnosis in patients with acute stress. 0 represented glioblastoma first symptoms, negative part represented time from stress event to first symptoms and positive part represented time from first symptoms to glioblastoma diagnosis.

	High stressors / High stress	High stressors / Low stress	Low stressors / Low stress	Low stressors / High stress
Number of patients	13 (20,3%)	8 (12,5%)	36 (56,3%)	7 (10,9%)
Median age at diagnosis (years)	58,5	63,4	69,3	66,7
Depression history	N = 8 (61,5%)	N = 4 (50%)	N = 4 (11,1%)	N = 3 (42,9%)
Psychotropic medication	N = 1 (7,7%)	N = 1 (12,5%)	N = 6 (16,7%)	N = 4 (57,1%)
Beta-blocker medication	N = 1 (7,7%)	N = 1 (12,5%)	N = 7 (19,4%)	N = 2 (28,6%)
Workers	N = 10 (76,9%)	N = 4 (50%)	N = 16 (44,4%)	N = 1 (14,3%)
Symptoms at diagnostic :				
- Deficit	N = 5 (38,5%)	N = 4 (50%)	N = 20 (55,6%)	N = 4 (57,1%)
- Cognitive impairment	N = 9 (69,2%)	N = 4 (50%)	N = 16 (44,4%)	N = 2 (28,6%)
- Epilepsy	N = 5 (38,5%)	N = 4 (50%)	N = 14 (38,9%)	N = 3 (42,3%)
- Intracranial hypertension	N = 0 (0%)	N = 1 (12,5%)	N = 1 (2,8%)	N = 1 (14,3%)
PS 0/1	N = 8 (61,5%)	N = 7 (87,5%)	N = 25 (69,4%)	N = 5 (71,4%)
Tumour location :				
- Frontal	N = 8 (61,5%)	N = 3 (37,5%)	N = 12 (33,3%)	N = 2 (28,6%)
- Temporal	N = 3 (23,1%)	N = 2 (25%)	N = 9 (25%)	N = 1 (14,3%)
- Parietal	N = 0	N = 0 (0%)	N = 7 (19,4%)	N = 1 (14,3%)
- Occipital	N = 0	N = 1 (12,5%)	N = 2 (5,6%)	N = 0
- Corpus callosum	N = 0	N = 1 (12,5%)	N = 1 (2,8%)	N = 0
- Thalamus	N = 0	N = 0	N = 0	N = 1 (14,3%)
- Multifocal	N = 2 (15,4%)	N = 1 (12,5%)	N = 5 (13,9%)	N = 2 (28,6%)
MGMT methylation	N = 3 (23,1%)	N = 1 (12,5%)	N = 14 (38,9%)	N = 2 (28,6%)
Median Time (month)				
- Acute stress - diagnosis	17	14	6	6

- First symptom - diagnosis	1.4	1.1	0.8	0.7
Psychological traumas	N = 11 (84,6%)	N = 7 (87,5%)	N = 9 (25%)	N = 4 (57,1%)

Table 1: Patients characteristics in the study population of IDH wild type glioblastoma patients.

Discussion

The present study described for the first time stressor exposure for patients with glioblastoma at diagnosis and stress response in this population. A majority of patients did not report acute stress prior to diagnosis of glioblastoma nor a high level of stress. PFS and OS were comparable in the different categories of stress response. The only factor that significantly and negatively affected PFS and OS in this study was the multifocal location.

Although stress represents an ambivalent response that can be beneficial in certain situations (concept of "fight or flight response"), the association between stress and pathologies are numerous and concern almost all medical specialties (depression, metabolic syndrome, coronary artery disease, psoriasis, inflammatory bowel disease etc.) [13-18]. Most epidemiological studies affirm that the prolonged perception of a high level of stress remains associated with reduced physical and mental condition [19,20]. This data is consistent with our results which find the higher rate of depression in the "high stressors / high stress" category of patients.

The autonomic nervous system and the hypothalamic-pituitary-adrenal system are thought to be responsible for the majority of interactions between stress and cancer via the secretion of catecholamines and cortisol [21-23]. We hypothesized that known inflammation and a decrease in immunoprotective functions induced by the secretion of cortisol and catecholamines could explain the mechanisms of gliomagenesis and glioma progression, already discussed in previous studies [21,24]. In several publications, stress-related psychosocial factors have been suggested to contribute to incidence of cancer, tumour progression and worst survival in cancer patients but most studies have focused on high-incidence cancer (especially in breast cancer [9,25-28]. Despite all the studies carried out, the stress-cancer association is currently not very robust. However, primary brain cancer incidence remains low (almost 20/100,000 person-years in France) and the risk factors for glioblastoma (brain radiation, genetic predisposition) are rarely found in medical practice [29]. The multiple questions from patients about the origin of the disease, led us to carry out this study and, to our knowledge, this is the first study to assess the impact of stress on the emergence of glioblastomas.

To address this question, we selected patients with primary brain tumours at the time of diagnosis to break free from stress induced by cancer management. Due to the Covid-19 health crisis during 2020, several eligible patients could not be included in our study. Furthermore, the study did not include cognitive test evaluation especially for patients excluded due to advanced cognitive disorder, incompatible with filling in the questionnaire. Moreover, we choose glioblastoma with a wild-type IDH profile, an aggressive tumour recognized for its rapid growth and patient symptoms briefly preceding diagnosis. This choice made it possible to limit the risk of patients presenting symptom-induced stress directly through the slow and chronic evolution of infiltrative gliomas with IDH mutation. [30]. Cardiovascular risk factors or coronaropathy represented a majority (56%) of medical histories of patients and these pathologies are known for their high-prevalence in the stressed population [6,31,32]. The depression rate found in this study (30%) was a little bit higher than in the general population (20%) and is probably explained by the cognitive impairment in the glioblastoma population [33]. The interaction of beta-blockers with the adrenergic pathway involved in stress and depression has been the subject of many publications concerning their impact on these pathologies and the proportion of patients treated with beta-blockers in our study seems similar to the proportion observed in the general population (20%) [34-38]. Location of the tumour did not impact stress, especially frontal location that can induce hypersensitivity which can lead to organic depression and anosodiaphoria which can lead to emotional indifference. The frontal lobe is involved in several functions including emotional regulation and a large portion of the patients in this study (up to 61,5% in the high stressor / high stress category) had an extensive frontal lobe tumour. In addition, these results are consistent because the frontal lobe represents the largest lobe of the brain and is one of the brain regions most affected by gliomas [39].

Some studies have described a better quality of life in cases of minor hemisphere syndrome [40]. The dominant hemisphere could be defined for all patients, however in our study there did not appear to be a significant difference in stress between left and

right hemisphere tumours. A large proportion of patients (56,3%) with recent diagnosis of glioblastoma reported no exposure to acute stress in the two years preceding the diagnosis. It seems important to emphasize that the "low stressor / low stress" patients corresponded to the oldest patients (pensioners). The proportion of job strain found in this study is low and the publications about job strain do not find a significant increase in risk of cancer [41,42]. Among the patients who identified recent acute stress, the time between the traumatic event and the first symptom was 11 months. In this single-center study with a socioculturally homogeneous population, health problems, loss of a loved one or job strain were the main stressors and certain types of stressors may have been underestimated (financial, healthcare access, war, etc.). Tumour progression and overall survival do not seem to be influenced by the existence or intensity of the patient's stress at diagnosis. However Liu, et al., demonstrated that additional stress in the history of patients due to poor prognostic and management are high, that we did not explore in the present study [43]. An important limit is the absence of a control cohort and therefore the lack of possibility of calculating the hazard ratio on stress exposure. In addition, the retrospective and declarative nature of the questionnaires may overestimate the stress level in the context of recent diagnosis of cancer. On the other hand, talking about stress and cancer could tend to relativize patient stress in comparison to the diagnosis of cancer and then minimize the presence of real stress that might have been important in another context. The cognitive distortion after receiving the diagnosis and the cognitive impairment induced by the tumour are probably responsible for a significant recall bias. Despite the inter and intra individual variability of stress as well as the multiple methodological difficulties to objectively evaluate this qualitative variable, several evaluation scales have been described for stressors and stress level and the Cungi scale used in our study has been validated and recognized as a sensitive and reproducible scale [12,43-47]. The stress questionnaire completed by patients was developed in 1997 by the French psychiatrist Charly Cungi, specialist in cognitive and behavioral therapy and author of numerous studies about stress. Thanks to the multiple dimensions explored as well as its sensitivity, reproducibility and reliability, this questionnaire constitutes a recognized and validated scale. Although some studies report a reduction in the risk of death from cancer, with no benefit in overall survival [21,48]. the personalized and multimodal management of the stress of our patients remains essential to guarantee the best possible quality of life. The survival benefit of stress management as well as the stress impact on efficacy of immunotherapy are other avenues worth exploring.

Conclusion

Although stress is an essential concern for patients with a primary brain tumour, psychological stress does not appear to impact the emergence of brain tumours nor the survival of patients with glioblastoma.

Conflict of Interest

The authors declared no potential conflicts of interest with respect to the research, authorship and/or publication of this article.

Funding Statement

This research did not receive any specific grant from funding agencies in the public, commercial or non-profit sectors.

Acknowledgement

None

Data Availability Statement

Not applicable.

Ethical Statement

The study was declared in CHU of Bordeaux ethic committee.

Informed Consent Statement

All patient agreed to participate and signed an informed consent.

Authors' Contributions

All authors contributed equally to this paper.

References

- Levav I, Rutz W. The WHO World Health Report 2001: new understanding: New hope. *ISR J Psychiatry Relat Sci.* 2002;39(1):50-6.
- Segerstrom SC, Miller GE. Psychological stress and the human immune system: A meta-analytic study of 30 years of inquiry. *Psychol Bull.* 2004;130(4):601-30.
- Rahe RH, Meyer M, Smith M, Kjaer G, Holmes TH. Social stress and illness onset. *J Psychosom Res.* 1964;8(1):35-44.
- Dhabhar FS. Effects of stress on immune function: the good, the bad and the beautiful. *Immunol Res.* 2014;58(2-3):193-210.
- Turner RJ, Wheaton B, Lloyd DA. The epidemiology of social stress. *Am Sociol Rev.* 1995;60(1):104.
- Steptoe A, Kivimäki M. Stress and cardiovascular disease: an update on current knowledge. *Annu Rev Public Health.* 2013;34(1):337-54.
- Lin Y, Wang C, Zhong Y. Striking life events associated with primary breast cancer susceptibility in women: A meta-analysis. *J Exp Clin Cancer Res.* 2013;32(1):53.
- Schoemaker MJ, Jones ME, Wright LB. Psychological stress, adverse life events and breast cancer incidence: A cohort investigation in 106,000 women in the United Kingdom. *Breast Cancer Res.* 2016;18(1):72.
- Chida Y, Hamer M, Wardle J, Steptoe A. Do stress-related psychosocial factors contribute to cancer incidence and survival? *Nat Rev Clin Oncol.* 2008;5(8):466-75.
- Ostrom QT, Bauchet L, Davis FG. The epidemiology of glioma in adults: A state-of-the-science review. *Neuro-Oncology.* 2014;16(7):896-913.
- Baldi I, Huchet A, Bauchet L, Loiseau H. Épidémiologie des glioblastomes. *Neurochirurgie.* 2010;56(6):433-40.
- Cungi C, Bouhana S, Degoul G, Bibollet D. Deux échelles brèves d'auto-évaluation des stresseurs et du stress. 1997.
- McEwen BS. Stress, adaptation and disease: Allostasis and allostatic load. *Ann N Y Acad Sci.* 1998;840(1):33-44.
- Hammen C. Stress and depression. *Annu Rev Clin Psychol.* 2005;1(1):293-319.
- Steptoe A, Kivimäki M. Stress and cardiovascular disease. *Nat Rev Cardiol.* 2012;9(6):360-70.
- Kimyai-Asadi A, Usman A. The role of psychological stress in skin disease. *J Cutan Med Surg.* 2001;5(2):140-5.
- Schoultz M, Beattie M, Gorely T, Leung J. Assessment of causal link between psychological factors and symptom exacerbation in inflammatory bowel disease. *Syst Rev.* 2020;9(1):169.
- Epel ES, Crosswell AD, Mayer SE. More than a feeling: A unified view of stress measurement for population science. *Front Neuroendocrinol.* 2018;49:146-169.
- Lantz PM, House JS, Mero RP, Williams DR. Stress, life events and socioeconomic disparities in health. *J Health Soc Behav.* 2005;46(3):274-88.
- Antoni MH, Dhabhar FS. The impact of psychosocial stress and stress management on immune responses in patients with cancer. *Cancer.* 2019;125(9):1417-31.
- Antoni MH, Lutgendorf SK, Cole SW. The influence of bio-behavioural factors on tumour biology. *Nat Rev Cancer.* 2006;6(3):240-8.
- Soygur H, Palaoglu O, Akarsu ES. Interleukin-6 levels and HPA axis activation in breast cancer patients with major depressive disorder. *Prog Neuropsychopharmacol Biol Psychiatry.* 2007;31(6):1242-7.
- Lu H. Inflammation, a key event in cancer development. *Mol Cancer Res.* 2006;4(4):221-33.
- Sloan EK, Priceman SJ, Cox BF. The sympathetic nervous system induces a metastatic switch in primary breast cancer. *Cancer Res.* 2010;70(18):7042-52.
- Antonova L, Aronson K, Mueller CR. Stress and breast cancer: From epidemiology to molecular biology. *Breast Cancer Res.* 2011;13(2):208.
- Duijts SFA, Zeegers MPA, van den Borne BV. Stressful life events and breast cancer risk: A meta-analysis. *Int J Cancer.* 2003;107(6):1023-9.
- Nielsen NR, Zhang ZF, Kristensen TS. Self-reported stress and risk of breast cancer. *BMJ.* 2005;331(7516):548.
- Pouchieu C, Gruber A, Berteaud E. Increasing incidence of CNS tumors (2000-2012). *BMC Cancer.* 2018;18(1):653.
- Bunevicius A, Deltuva VP, Tamasauskas A. Pre-operative depressive and anxiety symptoms and five-year survival of glioma and meningioma patients. *Oncotarget.* 2017;8(34):57543-51.
- Rozanski A, Blumenthal JA, Kaplan J. Psychological factors and cardiovascular disease. *Circulation.* 1999;99(16):2192-217.
- Kershaw KN, Brenes GA, Charles LE. Stressful life events and cardiovascular disease in women. *J Am Heart Assoc.* 2014;3(3).
- Fond G, Lancon C, Auquier P, Boyer L. Prevalence of major depression in France: A systematic review. *Presse Med.*

2019;48(4):365-75.

33. Chierichetti SM, Moise G, Galeone M. Beta-blockers and psychic stress. *Int J Clin Pharmacol Ther*. 1985;23(9):510-4.
34. Steptoe A, Ronaldson A, Kostich K. Beta-adrenergic blockade and stress responses. *Brain Behav Immun*. 2018;70:369-75.
35. Andrade C. β -Blockers and the risk of new-onset depression. *J Clin Psychiatry*. 2021;82(3):21f14095.
36. Riemer TG, Villagomez Fuentes LE, Algharably EAE. Do β -blockers cause depression? *Hypertension*. 2021;77(5):1539-48.
37. Bejan-Angoulvant T, Angoulvant D. Update on beta blockers in 2020. *Rev Med Interne*. 2020;41(11):741-7.
38. Larjavaara S, Mäntylä R, Salminen T. Incidence of gliomas by anatomic location. *Neuro Oncol*. 2007;9(3):319-25.
39. Lipsey JR, Robinson RG, Pearson GD. Mood change following bilateral hemisphere brain injury. *Br J Psychiatry*. 1983;143(3):266-73.
40. Vesterlund GK, Høeg BL, Johansen C. Prolonged job strain and cancer risk in women. *Acta Oncol*. 2017;56(2):301-6.
41. Heikkilä K, Nyberg ST, Theorell T. Work stress and cancer risk. *BMJ*. 2013;346:f165.
42. Liu F, Huang J, Zhang L. Screening for distress in patients with primary brain tumor. *BMC Cancer*. 2018;18(1):124.
43. Crosswell AD, Lockwood KG. Best practices for stress measurement. *Health Psychol Open*. 2020;7(2).
44. Slavich GM, Shields GS. Assessing lifetime stress exposure (Adult STRAIN). *Psychosom Med*. 2018;80(1):17-27.
45. Holmes TH, Rahe RH. The social readjustment rating scale. *J Psychosom Res*. 1967;11(2):213-8.
46. Lee EH. Review of the psychometric evidence of the perceived stress scale. *Asian Nurs Res*. 2012;6(4):121-7.
47. Oh PJ, Shin SR, Ahn HS, Kim HJ. Psychosocial interventions and cancer survival. *Psychol Health*. 2016;31(4):396-419.
48. Phillips KM, Antoni MH, Lechner SC. Stress management intervention in breast cancer. *Psychosom Med*. 2008;70(9):1044-9.

About the journal

Journal of Neuro and Oncology Research is an international, peer-reviewed, open-access journal published by Athenaeum Scientific Publishers. The journal publishes original research articles, case reports, editorials, reviews, and commentaries relevant to its scope. It aims to disseminate high-quality scholarly work that contributes to research, clinical practice, and academic knowledge in the field.

All submissions are evaluated through a structured peer-review process in accordance with established editorial and ethical standards. Manuscripts are submitted and processed through the journal's online submission system.

Manuscript submission: <https://athenaeumpub.com/submit-manuscript/>

